

NO DRAWINGS

- (21) Application No. 14529/69 (22) Filed 19 March 1969
 (23) Complete Specification filed 16 March 1970
 (45) Complete Specification published 20 Oct. 1971
 (51) International Classification C 07 d 29/28 29/20 99/02 57/00
 A 61 k 27/00//C 07 d 29/10 C 07 c 59/20 149/40
 (52) Index at acceptance

C2C 172—194—284 173—197—288 215 220 221 225 227
 22Y 246 247 250 251 25Y 28X 29X 29Y 304
 30Y 323 32Y 351 355 364 366 367 36Y 373 37Y
 3A12A4A 3A12A4C 3A12B3 3A12C1 3A12C6
 3A14A3D 3A14A5 3A14A8A 3A7V3A2 3A7V3E1
 3A7V3K1 456 45Y 463 464 552 553 612 613 61X
 620 625 628 660 662 665 668 675 770 771 790
 79Y LM NK QT QU ZD


- (72) Inventors OLE BENT TVAERMOSE NIELSEN, HANS-HASSO
 FREY and PETER WERNER FEIT

(54) PIPERIDINE COMPOUNDS FOR THE TREATMENT
 OF PARKINSONISM

(71) We, LØVENS KEMISKE FABRIK
 PRODUKTIONS-AKTIESELSKAB, a Company, in-
 5 corporated under the Laws of Denmark, of
 Ballerup, Denmark, do hereby declare the in-
 vention, for which we pray that a patent may
 be granted to us, and the method by which

it is to be performed, to be particularly
 described in and by the following state-
 ment:—

This invention relates to a series of hither- 10
 to unknown compounds of the general for-
 mula:



wherein R₁ is a straight or branched C3 to
 15 C12 aliphatic hydrocarbon chain, unsubsti-
 tuted or substituted with phenyl, phenoxy
 or phenylthio, which R₁ is directly attached
 to the benzene nucleus, or optionally may be
 20 attached to the benzene nucleus through a
 hetero atom which is an oxygen or sulphur
 atom; R₂ is alkyl, R₃ is a cycloalkyl radical
 with from 5 to 8 carbon atoms in the ring,
 and R₂ together with R₃ and the nitrogen atom
 25 can complete a heterocyclic ring which may
 be alkyl-substituted, n₁ is an integer from
 2 to 4, n₂ is an integer from 1 to 5, and
 n₃ is an integer from 1 to 3; to salts of the
 compounds with pharmaceutically acceptable
 30 inorganic and organic acids; and to methods
 for the preparation of the compounds and
 their salts.

When ever used in the statement above
 or in the description below, the term "alkyl"
 35 means lower-alkyl, including straight and
 branched aliphatic hydrocarbon chains with
 from 1 to 6 carbon atoms in the chain.

The compounds of the invention possess

valuable pharmacological activities, thus e.g.
 they display a favourable central anticholin- 40
 ergic action and are intended to be used in
 the treatment of patients suffering for in-
 stance from parkinsonism, including the post-
 encephalytic or arteriosclerotic parkinsonism
 and similar conditions.

As implied in the term, post-encephalytic 45
 parkinsonism refers to the appearance as a
 sequence to encephalitis of muscle rigidity
 and tremor frequently along with spasmodic
 phenomena, whereas the term arteriosclerotic
 parkinsonism refers to the appearance as a 50
 consequence of multiple cerebral vascular le-
 sions of difficulties of movements and fixity
 of posture, and similar conditions occurring
 in the older age group, often combined with
 muscle rigidity while tremor is absent. The 55
 said disorders are chronic and progressive and
 consequently all treatment is symptomatic
 and must be continued for long periods of
 time.

The medication may comprise treatment 60
 with belladonna alkaloids, e.g. atropine,

5 amphetamine alone or in combination with
belladonna alkaloids, with certain antihist-
aminics or apomorphine, and similar unspec-
ific medications, which may offer some degree
of symptomatic relief on tremor or spasmodic
conditions, but no fixed dosage can be recom-
mended and ordinarily small amounts of the
drug in question are used initially while larger
doses are ultimately required whereby it may
10 be necessary to approach the limit of toler-
ance and several toxic symptoms appear.
Better results in the treatment of parkin-
sonism have been observed by using certain
synthetic drugs as e.g. trihexaphenidyl (3 -
15 (1 - piperidyl) - 1 - phenyl - 1 - cyclo-
hexyl - 1 - propanol), Caramiphen (2 - di-
ethylaminoethyl - 1 - phenyl - cyclopentane -
1 - carboxylate hydrochloride), or Diethazide
20 (diethylaminoethyl - N - dibenzoparathia-
zine).

The action of trihexyphenidyl resembles
that of atropine, in particular as far as the
antispasmodic properties are concerned where-
as some of the undesired effects of atropine
25 are weaker, but still the peripheral para-
sympatholytic action of trihexyphenidyl must
be considered an undesired effect in the
treatment of parkinsonism where in particu-
lar the central action is important.

30 As far as the chemical constitution is
concerned the compounds of the present in-
vention differ far from the drugs mentioned
above, and it has surprisingly been found
that the compounds of formula I exert a
35 favourable specific therapeutic action with
a view to the treatment of all forms of
parkinsonism.

According to experiments the preferred
compounds with a view to treatment of par-
kinsonism are those of formula I in which
40 R_1 is a C5 to C7 aliphatic hydrocarbon
chain attached to the benzene nucleus through
the hetero atoms O or S, in which the integers
 n_1 and n_2 are within the limits from 2 to 3,
45 and from 2 to 4 respectively, and in which
 R_2 is a C1 to C2 aliphatic alkyl group, and
 R_3 is a C4 to C7 cycloalkyl group, or in
which R_2 and R_3 together with the N atom
form a heterocyclic ring.

50 In particular, however, the preferred com-
pounds are those in which R_1 has the meaning
defined above and are in the 4-position in
the benzene nucleus, and in which R_2 and
 R_3 together form an unsubstituted or or
55 alkyl-substituted pyrrolidino, piperidino,
hexamethylencimino or heptamethyleneimino
group.

Thus the compound 1 - (4 - n - hexyloxy-
benzoyl) - 4 - (piperidinoethyl) - piperidine
60 hydrochloride, among a series of related com-
pounds, displayed a promising central anti-
cholinergic activity, while its peripheral
parasympatholytic effects were comparably
weak. Its antagonistic effects against the
65 tremorogenic action of tremorine and oxotre-

morine, which is considered to be the most
predictive pharmacological model of parkin-
sonism, were two to five times stronger than
those of trihexyphenidyl being at present the
70 drug of choice in the treatment of parkin-
sonism. Furthermore, the central effects of
oxotremorine (tremor) were inhibited with
lower doses than the peripheral effects (sali-
vation) which as mentioned above is highly
75 desirable for antiparkinsonism drugs.

Experiments in higher animals further con-
firmed the favourable weak peripheral anti-
cholinergic action of the compounds of the
invention.

The acute oral toxicity of e.g. 1 - (4 - n -
80 hexyloxybenzoyl) - 4 - (piperidinoethyl) -
piperidine HCl expressed in LD₅₀ (mice)
is 470 mg/kg. which may be considered
low when compared to the degree of activity
85 in the anti-parkinson test in which an effect
could be observed with amounts of the order
0.5 to 2.0 mg/kg..

The chronic toxicity was studied in animal
experiments in which the test animals were
rats (Leo Wistar strain). The compounds were
90 administered orally each day in a period
of six months in various doses, in one animal
section in a daily dose of 50 mg/kg..

Even in this latter section no adverse
clinical signs were seen and no adverse changes
95 in bodyweight could be demonstrated. The
investigation comprises a full haematological
and pharmacological analysis of the animals
and after post-mortem examinations no ab-
normalities were demonstrated.

100 A pharmaceutical composition containing a
compound of the invention also constitutes
part of this invention. In the composition,
the proportion of therapeutically active
material to carrier substances and
105 auxiliary agents can vary between 0.04 to
10% depending upon the form of pharma-
ceutical presentation.

The composition in question can be worked
up to pharmaceutical forms of presentations
110 such as tablets, pills, dragees and supposi-
tories, or the composition can be filled in
medical containers such as capsules or am-
poules or, as far as mixtures or elixirs are
concerned, they may be filled in bottles
115 and similar containers.

Pharmaceutical inorganic or organic, solid
or liquid carriers suitable for enteral and
parantral administration can be used to make
up the composition; water, gelatine, lactose,
120 starch, magnesium stearate, talc, vegetable
and animal oils and fats, benzyl alcohol, gum,
polyalkylene glycol and similar other known
carriers for medicaments are suitable as car-
riers while stabilizing agents, wetting or
125 emulsifying agents, salts for varying the os-
motic pressure or buffers for securing an ade-
quate pH-value of the composition can be
used as auxiliary agents.

130 In the composition, the compounds of for-

of the compound of formula (II) may appropriately be employed in order to form e.g. the hydrogen halide of compounds of formula (I) directly in the reaction mixture.

- 5 In another embodiment the reaction is performed in the presence of an inert solvent, preferably immiscible with water, and at temperatures at or below 0° C. while the acid component possibly formed during the reaction e.g. a hydrogen halide, is continuously removed by adding an aqueous solution of a base, e.g. an alkali metal hydroxide. In this embodiment the starting substances are used in equivalent amounts, or substantially in equivalent amounts, and the reaction may be completed within a few hours.

- 10 After complete reaction, the desired compound is readily recovered from the organic phase, if necessary after having removed a possible excess of starting substance of formula (II) by extraction with an aqueous solution of an inorganic base, by evaporation of the organic phase, and recrystallizing the residue, or the compound may be isolated as a salt with an acid by neutralizing the base, in a suitable solvent or mixture of solvents with a view to the precipitation or crystallization of the salt.

- 15 The invention will now be illustrated by the following non-limiting Examples, of which Examples 1 to 4 illustrate the preparation of intermediates and Examples 5 to 8, illustrate the preparation of the compounds of formula (I):—

EXAMPLE 1

4-n-Hexylthiobenzoic acid

- 20 To a solution of 4 - aminobenzoic acid (32 g.), sodium nitrite (18.8 g.) and sodium hydroxide (11 g.) in water (150 ml.) concentrated hydrochloric acid (50 ml.) was slowly added while stirring rigorously at -5—0° C. After the addition was completed the stirring was continued for a further 1 hour at 0—5° C. The cooled diazonium-solution was filtered and slowly added to a solution of potassium xanthogenate (62.5 g.) and sodium carbonate (87.5 g.) in water (250 ml.) while stirring vigorously at 65—70° C. The mixture was stirred at 65—70° C. for a further 1 hour. After cooling the mixture was carefully acidified with concentrated hydrochloric acid (150 ml.). The precipitated material was filtered off, washed with water and dissolved in 10% sodium hydroxide solution (500 ml.). The flask was filled with nitrogen, closed and left overnight. n-Hexylbromide (85 g.) was added and the mixture was refluxed for 3 hours. The resulting mixture was poured into concentrated hydrochloric acid (200 ml.)/ice (about 200 g.), and the precipitate was filtered off and washed with water. After drying, 22 g. of crude 4-n-hexylthiobenzoic acid with a melting point of 89—93° C. was obtained. A sample repeatedly

recrystallized from cyclohexane had a melting point of 96—98° C. 65

EXAMPLE 2

4 - (4 - Phenylbutoxy) - benzoic acid

- A solution of ethyl 4-hydroxy benzoate (11 g.), 4 - phenylbutylbromide (17 g.) and sodium (1.53 g.) in ethanol (50 ml.) was refluxed for 20 hours and was then evaporated *in vacuo*. 4 N sodium hydroxide (25 ml.) was added to the residue, and the mixture was heated on a steam bath for 5 hours. After cooling the resulting solution was acidified with concentrated hydrochloric acid (15 ml.). The precipitated material was collected by filtration and washed with water. After drying, 4 - (4 - phenylbutoxy) - benzoic acid with a melting point of 128—131° C. was obtained. Recrystallization twice from aqueous ethanol raised the melting point to 130—132° C. By substituting in the above procedure equimolar amounts of 2 - n - butylthioethylchloride for the 4 - phenylbutylbromide, 4 - (2 - n - butylthioethoxy) - benzoic acid (m.p. 95—97° C.), was obtained. 70 75 80 85

EXAMPLE 3

4 - Piperidinomethyl - piperidine dihydrochloride hydrate

- To a stirred mixture of piperidine (12 g.), potassium carbonate (28 g.) and methanol (100 ml.), 4 - chloromethylpyridine hydrochloride (16.4 g.) was added in portions. The mixture was stirred at room temperature for a further 2 hours and was then evaporated *in vacuo*. The residue was treated with 2 N sodium hydroxide (25 ml.) and the separated oil was extracted with diethyl ether. The organic phase was dried (MgSO₄) and distilled. 4 - Piperidinomethylpyridine with a boiling point of 126—126.5° C. at 9 mm. Hg. was obtained. This material was dissolved in a mixture of methanol (75 ml.) and 3 N hydrochloric acid (45 ml.) and was hydrogenated after addition of PtO₂ (0.5 g.). The hydrogen uptake was complete within 3.5 hours. The catalyst was removed by filtration and the filtrate was evaporated *in vacuo*. The crystalline residue was triturated with acetone and collected by filtration. After drying, 4 - piperidinomethylpiperidine dihydrochloride hydrate with a melting point of about 260° C. was obtained. Recrystallization from ethanol raised the melting point to 265—266° C. 90 95 100 105 110 115

EXAMPLE 4

4 - [2 - (4 - Methylpiperidino) - ethyl] - piperidine dihydrochloride

- A mixture of 4-vinylpyridine (25 g.), 4 - methylpiperidine (35.4 g.) and acetic acid (3.5 ml.) was heated on a steam bath for 24 hours. 4 N Sodium hydroxide (25 ml.) was added to the cooled mixture and the separated oil was extracted with diethyl ether. 120 125

- The organic phase was dried (MgSO_4) and distilled. 4 - [2 - (4 - methylpiperidino) - ethyl] - pyridine with a boiling point of 151—154° C. at 9 mm. Hg. was obtained.
- 5 This material was dissolved in a mixture of methanol (230 ml.) and 4 N hydrochloric acid (130 ml.) and was hydrogenated after addition of PtO_2 (1.0 g.). The hydrogen uptake was complete within 20 hours. The catalyst was removed by filtration and the filtrate was evaporated *in vacuo*. The crystalline residue was triturated with acetone and was collected by filtration. After drying, 4 - [2 - (4 - methylpiperidino) - ethyl] - piperidine dihydrochloride with a melting point higher than 290° C. was obtained. By substituting in the above procedure equimolar amounts of 3 - methylpiperidine or N-methylcyclohexylamine for the 4 - methylpiperidine,
- 10 4 - [2 - (3 - methylpiperidino) - ethyl] - piperidine dihydrochloride (m.p. 267—269° C.) and 4 - (2 - N - methylcyclohexylaminoethyl) - piperidine dihydrochloride (hygroscopic) were obtained respectively.

EXAMPLE 5

- 1 - (4 - n - Hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride
- A solution of 4 - n - hexyloxybenzoyl chloride (5.0 g.) in methylenechloride (25 ml.) was slowly added to a mixture of 4 - (2 - piperidinoethyl) piperidine dihydrochloride (5.4 g.), methylenechloride (25 ml.) and 2 N sodium hydroxide (50 ml.) while stirring at 0—5° C. After the addition was complete the stirring was continued for a further 4 hours. The organic layer was separated, washed with brine, dried (MgSO_4) and evaporated *in vacuo*. The remaining material was dissolved in diethylether (50 ml.) and acidified with dry ethanolic hydrochloric acid. The precipitated oily material was crystallized from isopropanol/diethylether. After drying and recrystallization from acetone, 6.3 g. of 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride with a melting point of 197.5—198.5° C was obtained.

EXAMPLE 6

- By substituting in the above procedure equimolar amounts of 4 - n - propyloxybenzoyl chloride, 4 - isopropoxybenzoyl chloride, 4 - n - butyloxybenzoyl chloride, 4 - sec - butyloxybenzoyl chloride, 4 - isobutyloxybenzoyl chloride, 4 - isoamyloxybenzoyl chloride,
- 50 4 - n - heptyloxybenzoyl chloride, 4 - n - octyloxybenzoyl chloride, 3 - n - propyloxybenzoyl chloride, 3 - n - butyloxybenzoyl chloride, 3 - n - amyloxybenzoyl chloride, 3 - n - hexyloxybenzoyl chloride, 4 - n - hexylthiobenzoyle chloride, 4 - n - butylbenzoyl chloride, 4 - n - pentylbenzoyl chloride, 4 - n - hexylbenzoyl chloride, 4 - (2 - phenylethoxy) - benzoyl chloride, 4 - (3 -

- phenylpropoxy) - benzoyl chloride, 4 - (4 - phenyl butoxy) - benzoyl chloride, 4 - (2 - phenoxyethoxy) - benzoyl chloride, 4 - (2 - n - butylthioethoxy) - benzoyl chloride, 4 - n - heptylbenzoyl chloride, or 4 - n - octylbenzoyl chloride, for the 4 - n - hexyloxybenzoyl chloride, 1 - (4 - n - propyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 192—193.5° C.), 1 - (4 - isopropoxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 218—220° C.), 1 - (4 - n - butyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride, (m.p. 194.5—195.5° C.), 1 - (4 - sec. butyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 172—174° C.), 1 - (4 - isobutyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 189—191° C.), 1 - (4 - isoamyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 207—209° C.), 1 - (4 - n - heptyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 197—199° C.), 1 - (4 - n - octyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 198.5—199.5° C.), 1 - (3 - n - propyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride semihydrate (m.p. 156.5—158.5° C.), 1 - (3 - n - butyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 135.5—136.5° C.), 1 - (3 - n - amyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 124—126° C.), 1 - (3 - n - hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 135—136° C.), 1 - (4 - n - hexylthiobenzoyle) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 166—168° C.), 1 - (4 - n - butylbenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 171.5—172.5° C.), 1 - (4 - n - pentylbenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 170.5—172° C.), 1 - (4 - n - hexylbenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 172.5—173.5° C.),
- 1 - [4 - (2 - phenylethyl) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 190—191.5° C.), 1 - [4 - (3 - phenyl propoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 204—205° C.), 1 - [4 - (4 - phenylbutoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 158.5—160° C.), 1 - [4 - (2 - phenoxyethoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine (m.p. 103—106° C.), 1 - [4 - (2 - n - butylthioethoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 161—163° C.), 1 - (4 - n - heptylbenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 173—173.5° C.), and 1 (4 - n - octylbenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrochloride (m.p. 177—179° C.), were obtained.

piperidine hydrochloride (m.p. 178—180° C.) were obtained.

By substituting in the procedure described in the Example 5 above equimolar amounts of 4 - piperidinomethyl - piperidine dihydrochloride, 4 - (3 - piperidinopropyl) - piperidine dihydrochloride, 4 - (2 - N - methylcyclohexylaminoethyl)piperidine dihydrochloride, 4 - (2 - morpholinoethyl) - piperidine dihydrochloride, 3 - (piperidinomethyl) - piperidine dihydrochloride, 4 - (2 - pyrrolidinoethyl) - piperidine dihydrochloride, 4 - (2 - hexamethyleneimineethyl) - piperidine dihydrochloride, 4 - [2 - (3 - methylpiperidino) - ethyl] - piperidine dihydrochloride, or 4 - [2 - (4 - methylpiperidino) - ethyl] - piperidine hydrochloride, for the 4 - (2 - piperidinoethyl) - piperidine dihydrochloride, 1 - (4 - n - hexyloxybenzoyl) - 4 - piperidinomethyl piperidine hydrochloride (m.p. 199—200.5° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - (3 - piperidinopropyl) - piperidine hydrochloride (m.p. 147—149° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - N - methylcyclohexylaminoethyl) - piperidine dihydrochloride (m.p. 194—195° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - morpholinoethyl) - piperidine hydrochloride (m.p. 185.5—187° C.), 1 - (4 - n - hexyloxybenzoyl - 3 (piperidinomethyl) - piperidine hydrochloride (m.p. 149—151° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - pyrrolidinoethyl) - piperidine hydrochloride (m.p. 142.5—144.5° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - hexamethyleneimineethyl) - piperidine hydrochloride (m.p. 166.5—168.5° C.), 1 - (4 - n - hexyloxybenzoyl) - 4 - [2 - (3 - methylpiperidino) - ethyl] - piperidine hydrochloride (m.p. 174.5—177° C.), and 1 - (4 - n - hexyloxybenzoyl) - 4 - [2 - (4 - methyl - piperidino) - ethyl]

EXAMPLE 8

By using in the procedure described in the Example 5 above and substituting equimolar amounts of 4 - n - amyloxybenzoyl chloride for the 4 - n - hexyloxybenzoyl chloride and 48% hydrobromic acid for the ethanolic hydrochloric acid, 1 - (4 - n - amyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine hydrobromide (m.p. 192.5—194° C.) was obtained.

EXAMPLE 9

Tablets containing 5 mg. of the active compound, and the following composition, were prepared as described below:


1 - (4 - n - hexyloxybenzoyl) - 4 - (piperidinoethyl) - piperi- dine, hydrochloride	5 mg.	60
Lactose	69 mg.	
Starch	57 mg.	
Gelatin	2 mg.	
Talc	9 mg.	

The calculated amount of active compound, lactose, and starch, were mixed, granulated with a solution of gelatine in water and dried. After sifting the calculated amount of talc is added, the tablets are made by means of a 17 mm. punching die which provides tablets weighing 142 mg. each corresponding to 5 mg. of the active compound per tablet.

WHAT WE CLAIM IS:—

WHAT WE CLAIM IS:—

1. A compound of the general formula



wherein R_1 is a straight or branched C_3 to C_{12} aliphatic hydrocarbon chain, unsubstituted or substituted with phenyl, phenoxy or phenylthio, which R_1 is directly attached to the benzene nucleus, or optionally may be attached to the benzene nucleus through a hetero atom which is an oxygen or sulphur atom; R_2 is alkyl; R_3 is a cycloalkyl radical with from 5 to 8 carbon atoms in the ring, and R_2 together with R_3 can complete a heterocyclic ring which may be alkyl-substituted, n_1 is an integer from 2 to 4, n_2 is an integer from 1 to 5, and n_3 is an integer from 1 to 3; or a salt of the compound with a pharmaceutically acceptable inorganic or organic acid.

2. A compound according to Claim 1 in

which n_3 is 2 or 3, and R_1 is attached to the nucleus at the 4-position.

3. A compound according to Claims 1 and 2 in which n_1 and n_2 have the value 2, and R_1 is a C5 to C7 aliphatic hydrocarbon chain attached to the benzene nucleus through the hetero atom O.

4. A compound according to claims 1, 2 and 3 in which R₂ and R₃ form part of a heterocyclic ring.

5. A compound as claimed in claim 4, in which R₂ and R₃ form part of a heterocyclic ring selected from the group consisting of the pyrrolidino, the piperidino, the hexamethylenimino, and the heptamethylenimino rings.

6. A compound as claimed in claim 5 in 110

which R_1 is the n-pentyl or the n-hexyl group.

7. 1 - [4 - (3 - phenylpropoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine and salts thereof.

5 8. 1 - [4 - (2 - phenoxyethoxy) - benzoyl] - 4 - (2 - piperidinoethyl) - piperidine and salts thereof.



9. 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - hexamethyleneiminoethyl) - piperidine and salts thereof.

10 10. 1 - (4 - n - hexyloxybenzoyl) - 4 - [2 - (4 - methylpiperidino) - ethyl] - piperidine and salts thereof.

11. 1 - (4 - n - amyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine and salts thereof.

12. 1 - (4 - n - hexyloxybenzoyl) - 4 - (piperidinoethyl) - piperidine and salts thereof.

20 13. A method of producing a compound of the general formula I as claimed in Claim 1 wherein an acid of the general formula (II)




45 wherein R_1 is a straight or branched C4 to C12 aliphatic hydrocarbon chain, unsubstituted or substituted with phenyl, phenoxy or phenylthio, which R_1 is directly attached to the benzene nucleus, or optionally may be attached to the benzene nucleus through a hetero atom which is an oxygen or sulphur atom; R_2 is alkyl; R_3 is a cycloalkyl radical with from 5 to 8 carbon atoms in the ring, and R_2 together with R_3 can complete a heterocyclic ring which may be alkyl-substituted, n_1 is an integer from 2 to 4, n_2 is an integer from 1 to 5, and n_3 is an integer from 1 to 3; and its salts with pharmaceutically acceptable inorganic and organic acids together with an atoxic pharmaceutically acceptable carrier, the quantity of the said compound of formula I in the unit being between 0.1 and 50 mg. calculated as the free base.

65 15. A pharmaceutical preparation in oral dosage unit form as claimed in claim 14, in which the units contain from 0.5 to 25 mg. of 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine.

70 16. A pharmaceutical preparation in dosage form as claimed in claim 14 or 15,

in which R_1 has the meaning hereinbefore defined is reacted in the form on an acid halide, an anhydride, a mixed anhydride with an alkyl-carbonic acid, a carboxylic acid, a sulphonic acid or with an inorganic acid, or in the form of a reactive derivative obtained by reacting the acid with a carbodiimide or N',N' - carbonyldiimidazole, with a compound of the general formula:



in which R_2 , R_3 , n_1 , n_2 and n_3 have the meanings hereinbefore defined, whereafter the compound formed is recovered as such or as one of its salts with acids.

14. A pharmaceutical preparation in dosage unit form for the treatment of patients suffering from parkinsonism comprising as at least one active component a compound of the general formula I

in which the units contain from 1 to 10 mg. of 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine.

17. A pharmaceutical preparation in dosage unit form as claimed in any one of Claims 14 to 16, in the form of pills, tablets or capsules.

18. A pharmaceutical preparation in dosage unit form as claimed in Claim 14, in which the preparation is an injectable preparation containing from 0.1 to 25 mg. of the compound of formula I.

19. A pharmaceutical preparation in dosage unit form as claimed in Claim 18, in which the active ingredient is 1 - (4 - n - hexyloxybenzoyl) - 4 - (2 - piperidinoethyl) - piperidine in the form of one of its salts with a non-toxic acid dissolved in an aqueous medium.

20. A compound of the general formula 1 defined in Claim 1 substantially as hereinbefore described in any one of Examples 1 to 8 of the foregoing Examples.

21. A method of producing a compound of the general formula I defined in Claim 1 substantially as hereinbefore described in any

one of Examples 1 to 8 of the foregoing Examples.

22. A pharmaceutical preparation in dosage unit form substantially as hereinbefore described in Example 9 of the foregoing Examples.
- 5

TREGEAR, THIEMANN & BLEACH,
Chartered Patent Agents,
Melbourne House, Aldwych,
London, W.C.2.
Agents for the Applicant(s).

Printed for Her Majesty's Stationery Office, by the Courier Press, Leamington Spa, 1971.
Published by The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.